

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, C07K 14/47, C12N 15/62, C07K 16/18	A2	(11) International Publication Number: WO 00/15796 (43) International Publication Date: 23 March 2000 (23.03.00)
(21) International Application Number: PCT/US99/21090 (22) International Filing Date: 15 September 1999 (15.09.99) (30) Priority Data: PCT/US98/19330 16 September 1998 (16.09.98) US (71) Applicant (for all designated States except US): GENENTECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CHEN, Jian [CN/US]; 121 York Drive, Princeton, NJ 08540 (US). GODDARD, Audrey [CA/US]; 110 Congo Street, San Francisco, CA 94131 (US). GURNEY, Austin, L. [US/US]; 1 Debbie Lane, Belmont, CA 94002 (US). HILLAN, Kenneth [GB/US]; 64 Seward Street, San Francisco, CA 94114 (US). PENNICA, Diane [US/US]; 2417 Hale Drive, Burlingame, CA 94010 (US). WOOD, William, I. [US/US]; 35 Southdown Court, Hillsborough, CA 94010 (US). YUAN, Jean [CN/US]; 176 West 37th Avenue, San Mateo, CA 94003 (US). (74) Agents: KRESNAK, Mark, T. et al.; Genentech Inc., 1 DNA Way, South San Francisco, CA 94080-4990 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC ACIDS ENCODING THE SAME		
(57) Abstract The present invention is directed to novel polypeptides and to nucleic acid molecules encoding those polypeptides. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention.		

nucleotides

46. **PRO268**

Protein disulfide isomerase is an enzymatic protein which is involved in the promotion of correct refolding of proteins through the establishment of correct disulfide bond formation. Protein disulfide isomerase was initially identified based upon its ability to catalyze the renaturation of reduced denatured RNase (Goldberger et al., *J. Biol. Chem.* 239:1406-1410 (1964) and Epstein et al., *Cold Spring Harbor Symp. Quant. Biol.* 28:439-449 (1963)). Protein disulfide isomerase has been shown to be a resident enzyme of the endoplasmic reticulum which is retained in the endoplasmic reticulum via a -KDEL or -HDEL amino acid sequence at its C-terminus.

Given the importance of disulfide bond-forming enzymes and their potential uses in a number of different applications, for example in increasing the yield of correct refolding of recombinantly produced proteins, efforts are currently being undertaken by both industry and academia to identify new, native proteins having homology to protein disulfide isomerase. Many of these efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel protein disulfide isomerase homologs. We herein describe a novel polypeptide having homology to protein disulfide isomerase, designated herein as PRO268.

47. **PRO330**

Prolyl 4-hydroxylase is an enzyme which functions to post-translationally hydroxylate proline residues at the Y position of the amino acid sequence Gly-X-Y, which is a repeating three amino acid sequence found in both collagen and procollagen. Hydroxylation of proline residues at the Y position of the Gly-X-Y amino acid triplet to form 4-hydroxyproline residues at those positions is required before newly synthesized collagen polypeptide chains may fold into their proper three-dimensional triple-helical conformation. If hydroxylation does not occur, synthesized collagen polypeptides remain non-helical, are poorly secreted by cells and cannot assemble into stable functional collagen fibrils. Vuorio et al., *Proc. Natl. Acad. Sci. USA* 89:7467-7470 (1992). Prolyl 4-hydroxylase is comprised of at least two different polypeptide subunits, alpha and beta.

Efforts are being undertaken by both industry and academia to identify new, native secreted and membrane-bound receptor proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted and membrane-bound receptor proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.*, 93:7108-7113 (1996); U.S. Patent No. 5,536,637]. Based upon these efforts, Applicants have herein identified and describe a novel polypeptide having homology to the alpha subunit of prolyl 4-hydroxylase, designated herein as PRO330.

48. **PRO339 and PRO310**

Fringe is a protein which specifically blocks serrate-mediated activation of notch in the dorsal compartment of the *Drosophila* wing imaginal disc. Fleming, et al., *Development*, 124(15):2973-81 (1997).

Therefore, fringe is of interest for both its role in development as well as its ability to regulate serrate, particularly serrate's signaling abilities. Also of interest are novel polypeptides which may have a role in development and/or the regulation of serrate-like molecules. Of particular interest are novel polypeptides having homology to fringe as identified and described herein, designated herein as PRO339 and PRO310 polypeptides.

5 49. **PRO244**

Lectins are a class of proteins comprising a region that binds carbohydrates specifically and non-covalently. Numerous lectins have been identified in higher animals, both membrane-bound and soluble, and have been implicated in a variety of cell-recognition phenomena and tumor metastasis.

Most lectins can be classified as either C-type (calcium-dependent) or S-type (thiol-dependent).

10 Lectins are thought to play a role in regulating cellular events that are initiated at the level of the plasma membrane. For example, plasma membrane associated molecules are involved in the activation of various subsets of lymphoid cells, e.g. T-lymphocytes, and it is known that cell surface molecules are responsible for activation of these cells and consequently their response during an immune reaction.

A particular group of cell adhesion molecules, selectins, belong in the superfamily of C-type lectins.

15 This group includes L-selectin (peripheral lymph node homing receptor (pnHR), LEC-CAM-1, LAM-1, gp90^{MEL}, gp100^{MEL}, gp110^{MEL}, MEL-14 antigen, Leu-8 antigen, TQ-1 antigen, DREG antigen), E-selectin (LEC-CAM-2, LECAM-2, ELAM-1), and P-selectin (LEC-CAM-3, LECAM-3, GMP-140, PADGEM). The structure of selectins consists of a C-type lectin (carbohydrate binding) domain, an epidermal growth factor-like (EGF-like) motif, and variable numbers of complement regulatory (CR) motifs. Selectins are associated with
20 leukocyte adhesion, e.g. the attachment of neutrophils to venular endothelial cells adjacent to inflammation (E-selectin), or with the trafficking of lymphocytes from blood to secondary lymphoid organs, e.g. lymph nodes and Peyer's patches (L-selectin).

Another exemplary lectin is the cell-associated macrophage antigen, Mac-2 that is believed to be involved in cell adhesion and immune responses. Macrophages also express a lectin that recognizes Tn Ag, a
25 human carcinoma-associated epitope.

Another C-type lectin is CD95 (Fas antigen/APO-1) that is an important mediator of immunologically relevant regulated or programmed cell death (apoptosis). "Apoptosis" is a non-necrotic cell death that takes place in metazoan animal cells following activation of an intrinsic cell suicide program. The cloning of Fas antigen is described in PCT publication WO 91/10448, and European patent application EP510691. The mature
30 Fas molecule consists of 319 amino acids of which 157 are extracellular, 17 constitute the transmembrane domain, and 145 are intracellular. Increased levels of Fas expression at T cell surface have been associated with tumor cells and HIV-infected cells. Ligation of CD95 triggers apoptosis in the presence of interleukin-1 (IL-2).

C-type lectins also include receptors for oxidized low-density lipoprotein (LDL). This suggests a possible role in the pathogenesis of atherosclerosis.

35 We herein describe the identification and characterization of novel polypeptides having homology to C-type lectins, designated herein as PRO244 polypeptides.

In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO330 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO330 polypeptide having amino acid residues 1 to 533 of Figure 116 (SEQ ID NO:332), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

5 In another embodiment, the invention provides isolated PRO330 polypeptide. In particular, the invention provides isolated native sequence PRO330 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 533 of Figure 116 (SEQ ID NO:332).

48. **PRO339 and PRO310**

10 Applicants have identified two cDNA clones wherein each clone encodes a novel polypeptide having homology to fringe, wherein the polypeptides are designated in the present application as "PRO339" and "PRO310".

In one embodiment, the invention provides isolated nucleic acid molecules comprising DNA encoding a PRO339 and/or a PRO310 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the
15 PRO339 polypeptide having amino acid residues 1 to 772 of Figure 118 (SEQ ID NO:339), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions. In another aspect, the isolated nucleic acid comprises DNA encoding the PRO310 polypeptide having amino acid residues 1 to 318 of Figure 120 (SEQ ID NO:341), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally,
20 under high stringency conditions.

In another embodiment, the invention provides isolated PRO339 as well as isolated PRO310 polypeptides. In particular, the invention provides isolated native sequence PRO339 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 772 of Figure 118 (SEQ ID NO:339). The invention further provides isolated native sequence PRO310 polypeptide, which in one embodiment, includes
25 an amino acid sequence comprising residues 1 to 318 of Figure 120 (SEQ ID NO:341).

49. **PRO244**

Applicants have identified a cDNA clone that encodes a novel polypeptide, designated in the present application as "PRO244".

30 In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding PRO244 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding PRO244 polypeptide having amino acid residues 1 to 219 of Fig. 122 (SEQ ID NO:377), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

35 In another embodiment, the invention provides isolated PRO244 polypeptide. In particular, the invention provides isolated native sequence PRO244 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 219 of Figure 122 (SEQ ID NO:377).

50. Additional Embodiments

In other embodiments of the present invention, the invention provides vectors comprising DNA encoding any of the herein described polypeptides. Host cell comprising any such vector are also provided. By way of example, the host cells may be CHO cells, *E. coli*, or yeast. A process for producing any of the herein described polypeptides is further provided and comprises culturing host cells under conditions suitable for
5 expression of the desired polypeptide and recovering the desired polypeptide from the cell culture.

In other embodiments, the invention provides chimeric molecules comprising any of the herein described polypeptides fused to a heterologous polypeptide or amino acid sequence. Example of such chimeric molecules comprise any of the herein described polypeptides fused to an epitope tag sequence or a Fc region of an immunoglobulin.

10 In another embodiment, the invention provides an antibody which specifically binds to any of the above or below described polypeptides. Optionally, the antibody is a monoclonal antibody, humanized antibody, antibody fragment or single-chain antibody.

In yet other embodiments, the invention provides oligonucleotide probes useful for isolating genomic and cDNA nucleotide sequences, wherein those probes may be derived from any of the above or below described
15 nucleotide sequences.

In other embodiments, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a PRO polypeptide.

In one aspect, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82%
20 sequence identity, yet more preferably at least about 83% sequence identity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity, yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at least about 88% sequence identity, yet more preferably at least about 89% sequence identity, yet more preferably at least about 90% sequence identity, yet more preferably at least about 91% sequence identity, yet more
25 preferably at least about 92% sequence identity, yet more preferably at least about 93% sequence identity, yet more preferably at least about 94% sequence identity, yet more preferably at least about 95% sequence identity, yet more preferably at least about 96% sequence identity, yet more preferably at least about 97% sequence identity, yet more preferably at least about 98% sequence identity and yet more preferably at least about 99% sequence identity to (a) a DNA molecule encoding a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein or an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein, or (b) the complement of the DNA molecule of (a).

In other aspects, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82%
35 sequence identity, yet more preferably at least about 83% sequence identity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity, yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at

least about 88 % sequence identity, yet more preferably at least about 89 % sequence identity, yet more preferably at least about 90 % sequence identity, yet more preferably at least about 91 % sequence identity, yet more preferably at least about 92 % sequence identity, yet more preferably at least about 93 % sequence identity, yet more preferably at least about 94 % sequence identity, yet more preferably at least about 95 % sequence identity, yet more preferably at least about 96 % sequence identity, yet more preferably at least about 97 % sequence identity, yet more preferably at least about 98 % sequence identity and yet more preferably at least about 99 % sequence identity to (a) a DNA molecule comprising the coding sequence of a full-length PRO polypeptide cDNA as disclosed herein, the coding sequence of a PRO polypeptide lacking the signal peptide as disclosed herein or the coding sequence of an extracellular domain of a transmembrane PRO polypeptide, with or without the signal peptide, as disclosed herein, or (b) the complement of the DNA molecule of (a).

In a further aspect, the invention concerns an isolated nucleic acid molecule comprising a nucleotide sequence having at least about 80 % sequence identity, preferably at least about 81 % sequence identity, more preferably at least about 82 % sequence identity, yet more preferably at least about 83 % sequence identity, yet more preferably at least about 84 % sequence identity, yet more preferably at least about 85 % sequence identity, yet more preferably at least about 86 % sequence identity, yet more preferably at least about 87 % sequence identity, yet more preferably at least about 88 % sequence identity, yet more preferably at least about 89 % sequence identity, yet more preferably at least about 90 % sequence identity, yet more preferably at least about 91 % sequence identity, yet more preferably at least about 92 % sequence identity, yet more preferably at least about 93 % sequence identity, yet more preferably at least about 94 % sequence identity, yet more preferably at least about 95 % sequence identity, yet more preferably at least about 96 % sequence identity, yet more preferably at least about 97 % sequence identity, yet more preferably at least about 98 % sequence identity and yet more preferably at least about 99 % sequence identity to (a) a DNA molecule that encodes the same mature polypeptide encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein, or (b) the complement of the DNA molecule of (a).

Another aspect the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding a PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated, or is complementary to such encoding nucleotide sequence, wherein the transmembrane domain(s) of such polypeptide are disclosed herein. Therefore, soluble extracellular domains of the herein described PRO polypeptides are contemplated.

Another embodiment is directed to fragments of a PRO polypeptide coding sequence, or the complement thereof, that may find use as, for example, hybridization probes or for encoding fragments of a PRO polypeptide that may optionally encode a polypeptide comprising a binding site for an anti-PRO antibody. Such nucleic acid fragments are usually at least about 20 nucleotides in length, preferably at least about 30 nucleotides in length, more preferably at least about 40 nucleotides in length, yet more preferably at least about 50 nucleotides in length, yet more preferably at least about 60 nucleotides in length, yet more preferably at least about 70 nucleotides in length, yet more preferably at least about 80 nucleotides in length, yet more preferably at least about 90 nucleotides in length, yet more preferably at least about 100 nucleotides in length, yet more preferably at least about 110 nucleotides in length, yet more preferably at least about 120 nucleotides in length, yet more

preferably at least about 130 nucleotides in length, yet more preferably at least about 140 nucleotides in length, yet more preferably at least about 150 nucleotides in length, yet more preferably at least about 160 nucleotides in length, yet more preferably at least about 170 nucleotides in length, yet more preferably at least about 180 nucleotides in length, yet more preferably at least about 190 nucleotides in length, yet more preferably at least about 200 nucleotides in length, yet more preferably at least about 250 nucleotides in length, yet more preferably at least about 300 nucleotides in length, yet more preferably at least about 350 nucleotides in length, yet more preferably at least about 400 nucleotides in length, yet more preferably at least about 450 nucleotides in length, yet more preferably at least about 500 nucleotides in length, yet more preferably at least about 600 nucleotides in length, yet more preferably at least about 700 nucleotides in length, yet more preferably at least about 800 nucleotides in length, yet more preferably at least about 900 nucleotides in length and yet more preferably at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. It is noted that novel fragments of a PRO polypeptide-encoding nucleotide sequence may be determined in a routine manner by aligning the PRO polypeptide-encoding nucleotide sequence with other known nucleotide sequences using any of a number of well known sequence alignment programs and determining which PRO polypeptide-encoding nucleotide sequence fragment(s) are novel. All of such PRO polypeptide-encoding nucleotide sequences are contemplated herein. Also contemplated are the PRO polypeptide fragments encoded by these nucleotide molecule fragments, preferably those PRO polypeptide fragments that comprise a binding site for an anti-PRO antibody.

In another embodiment, the invention provides isolated PRO polypeptide encoded by any of the isolated nucleic acid sequences hereinabove identified.

In a certain aspect, the invention concerns an isolated PRO polypeptide, comprising an amino acid sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82% sequence identity, yet more preferably at least about 83% sequence identity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity, yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at least about 88% sequence identity, yet more preferably at least about 89% sequence identity, yet more preferably at least about 90% sequence identity, yet more preferably at least about 91% sequence identity, yet more preferably at least about 92% sequence identity, yet more preferably at least about 93% sequence identity, yet more preferably at least about 94% sequence identity, yet more preferably at least about 95% sequence identity, yet more preferably at least about 96% sequence identity, yet more preferably at least about 97% sequence identity, yet more preferably at least about 98% sequence identity and yet more preferably at least about 99% sequence identity to a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein or an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein.

In a further aspect, the invention concerns an isolated PRO polypeptide comprising an amino acid sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82% sequence identity, yet more preferably at least about 83% sequence identity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity,

yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at least about 88% sequence identity, yet more preferably at least about 89% sequence identity, yet more preferably at least about 90% sequence identity, yet more preferably at least about 91% sequence identity, yet more preferably at least about 92% sequence identity, yet more preferably at least about 93% sequence identity, yet more preferably at least about 94% sequence identity, yet more preferably at least about 95% sequence identity, yet more preferably at least about 96% sequence identity, yet more preferably at least about 97% sequence identity, yet more preferably at least about 98% sequence identity and yet more preferably at least about 99% sequence identity to an amino acid sequence encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein.

In a further aspect, the invention concerns an isolated PRO polypeptide comprising an amino acid sequence scoring at least about 80% positives, preferably at least about 81% positives, more preferably at least about 82% positives, yet more preferably at least about 83% positives, yet more preferably at least about 84% positives, yet more preferably at least about 85% positives, yet more preferably at least about 86% positives, yet more preferably at least about 87% positives, yet more preferably at least about 88% positives, yet more preferably at least about 89% positives, yet more preferably at least about 90% positives, yet more preferably at least about 91% positives, yet more preferably at least about 92% positives, yet more preferably at least about 93% positives, yet more preferably at least about 94% positives, yet more preferably at least about 95% positives, yet more preferably at least about 96% positives, yet more preferably at least about 97% positives, yet more preferably at least about 98% positives and yet more preferably at least about 99% positives when compared with the amino acid sequence of a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein or an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein.

In a specific aspect, the invention provides an isolated PRO polypeptide without the N-terminal signal sequence and/or the initiating methionine and is encoded by a nucleotide sequence that encodes such an amino acid sequence as hereinbefore described. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

Another aspect the invention provides an isolated PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

In yet another embodiment, the invention concerns agonists and antagonists of a native PRO polypeptide as defined herein. In a particular embodiment, the agonist or antagonist is an anti-PRO antibody or a small molecule.

In a further embodiment, the invention concerns a method of identifying agonists or antagonists to a PRO polypeptide which comprise contacting the PRO polypeptide with a candidate molecule and monitoring a

biological activity mediated by said PRO polypeptide. Preferably, the PRO polypeptide is a native PRO polypeptide.

In a still further embodiment, the invention concerns a composition of matter comprising a PRO polypeptide, or an agonist or antagonist of a PRO polypeptide as herein described, or an anti-PRO antibody, in combination with a carrier. Optionally, the carrier is a pharmaceutically acceptable carrier.

- 5 Another embodiment of the present invention is directed to the use of a PRO polypeptide, or an agonist or antagonist thereof as hereinbefore described, or an anti-PRO antibody, for the preparation of a medicament useful in the treatment of a condition which is responsive to the PRO polypeptide, an agonist or antagonist thereof or an anti-PRO antibody.

10 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a nucleotide sequence (SEQ ID NO:1) of a native sequence PRO211 cDNA, wherein SEQ ID NO:1 is a clone designated herein as "DNA32292-1131".

Figure 2 shows the amino acid sequence (SEQ ID NO:2) derived from the coding sequence of SEQ ID NO:1 shown in Figure 1.

- 15 Figure 3 shows a nucleotide sequence (SEQ ID NO:3) of a native sequence PRO217 cDNA, wherein SEQ ID NO:3 is a clone designated herein as "DNA33094-1131".

Figure 4 shows the amino acid sequence (SEQ ID NO:4) derived from the coding sequence of SEQ ID NO:3 shown in Figure 3.

- 20 Figure 5 shows a nucleotide sequence (SEQ ID NO:11) of a native sequence PRO230 cDNA, wherein SEQ ID NO:11 is a clone designated herein as "DNA33223-1136".

Figure 6 shows the amino acid sequence (SEQ ID NO:12) derived from the coding sequence of SEQ ID NO:11 shown in Figure 5.

Figure 7 shows a nucleotide sequence designated herein as DNA20088 (SEQ ID NO:13).

- 25 Figure 8 shows a nucleotide sequence (SEQ ID NO:17) of a native sequence PRO232 cDNA, wherein SEQ ID NO:17 is a clone designated herein as "DNA34435-1140".

Figure 9 shows the amino acid sequence (SEQ ID NO:18) derived from the coding sequence of SEQ ID NO:17 shown in Figure 8.

Figure 10 shows a nucleotide sequence (SEQ ID NO:22) of a native sequence PRO187 cDNA, wherein SEQ ID NO:22 is a clone designated herein as "DNA27864-1155".

- 30 Figure 11 shows the amino acid sequence (SEQ ID NO:23) derived from the coding sequence of SEQ ID NO:22 shown in Figure 10.

Figure 12 shows a nucleotide sequence (SEQ ID NO:27) of a native sequence PRO265 cDNA, wherein SEQ ID NO:27 is a clone designated herein as "DNA36350-1158".

- 35 Figure 13 shows the amino acid sequence (SEQ ID NO:28) derived from the coding sequence of SEQ ID NO:27 shown in Figure 12.

Figure 14 shows a nucleotide sequence (SEQ ID NO:33) of a native sequence PRO219 cDNA, wherein SEQ ID NO:33 is a clone designated herein as "DNA32290-1164".

Figure 110 shows the amino acid sequence (SEQ ID NO:315) derived from the coding sequence of SEQ ID NO:314 shown in Figure 109.

Figure 111 shows a nucleotide sequence (SEQ ID NO:319) of a native sequence PRO346 cDNA, wherein SEQ ID NO:319 is a clone designated herein as "DNA44167-1243".

5 Figure 112 shows the amino acid sequence (SEQ ID NO:320) derived from the coding sequence of SEQ ID NO:319 shown in Figure 111.

Figure 113 shows a nucleotide sequence (SEQ ID NO:324) of a native sequence PRO268 cDNA, wherein SEQ ID NO:324 is a clone designated herein as "DNA39427-1179".

Figure 114 shows the amino acid sequence (SEQ ID NO:325) derived from the coding sequence of SEQ ID NO:324 shown in Figure 113.

10 Figure 115 shows a nucleotide sequence (SEQ ID NO:331) of a native sequence PRO330 cDNA, wherein SEQ ID NO:331 is a clone designated herein as "DNA40603-1232".

Figure 116 shows the amino acid sequence (SEQ ID NO:332) derived from the coding sequence of SEQ ID NO:331 shown in Figure 115.

15 Figure 117 shows a nucleotide sequence (SEQ ID NO:338) of a native sequence PRO339 cDNA, wherein SEQ ID NO:338 is a clone designated herein as "DNA43466-1225".

Figure 118 shows the amino acid sequence (SEQ ID NO:339) derived from the coding sequence of SEQ ID NO:338 shown in Figure 117.

Figure 119 shows a nucleotide sequence (SEQ ID NO:340) of a native sequence PRO310 cDNA, wherein SEQ ID NO:340 is a clone designated herein as "DNA43046-1225".

20 Figure 120 shows the amino acid sequence (SEQ ID NO:341) derived from the coding sequence of SEQ ID NO:340 shown in Figure 119.

Figure 121 shows a nucleotide sequence (SEQ ID NO:376) of a native sequence PRO244 cDNA, wherein SEQ ID NO:376 is a clone designated herein as "DNA35668-1171".

25 Figure 122 shows the amino acid sequence (SEQ ID NO:377) derived from the coding sequence of SEQ ID NO:376 shown in Figure 121.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTSI. Definitions

The terms "PRO polypeptide" and "PRO" as used herein and when immediately followed by a numerical designation refer to various polypeptides, wherein the complete designation (i.e., PRO/number) refers to specific polypeptide sequences as described herein. The terms "PRO/number polypeptide" and "PRO/number" wherein the term "number" is provided as an actual numerical designation as used herein encompass native sequence polypeptides and polypeptide variants (which are further defined herein). The PRO polypeptides described herein may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods.

A "native sequence PRO polypeptide" comprises a polypeptide having the same amino acid sequence as the corresponding PRO polypeptide derived from nature. Such native sequence PRO polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. The term "native sequence PRO polypeptide" specifically encompasses naturally-occurring truncated or secreted forms of the specific PRO polypeptide (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. In various embodiments of the invention, the native sequence PRO polypeptides disclosed herein are mature or full-length native sequence polypeptides comprising the full-length amino acids sequences shown in the accompanying figures. Start and stop codons are shown in bold font and underlined in the figures. However, while the PRO polypeptide disclosed in the accompanying figures are shown to begin with methionine residues designated herein as amino acid position 1 in the figures, it is conceivable and possible that other methionine residues located either upstream or downstream from the amino acid position 1 in the figures may be employed as the starting amino acid residue for the PRO polypeptides.

The PRO polypeptide "extracellular domain" or "ECD" refers to a form of the PRO polypeptide which is essentially free of the transmembrane and cytoplasmic domains. Ordinarily, a PRO polypeptide ECD will have less than 1% of such transmembrane and/or cytoplasmic domains and preferably, will have less than 0.5% of such domains. It will be understood that any transmembrane domains identified for the PRO polypeptides of the present invention are identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than about 5 amino acids at either end of the domain as initially identified herein. Optionally, therefore, an extracellular domain of a PRO polypeptide may contain from about 5 or fewer amino acids on either side of the transmembrane domain/extracellular domain boundary as identified in the Examples or specification and such polypeptides, with or without the associated signal peptide, and nucleic acid encoding them, are contemplated by the present invention.

The approximate location of the "signal peptides" of the various PRO polypeptides disclosed herein are shown in the accompanying figures. It is noted, however, that the C-terminal boundary of a signal peptide may vary, but most likely by no more than about 5 amino acids on either side of the signal peptide C-terminal boundary as initially identified herein, wherein the C-terminal boundary of the signal peptide may be identified pursuant to criteria routinely employed in the art for identifying that type of amino acid sequence element (e.g.,

Nielsen et al., Prot. Eng. 10:1-6 (1997) and von Heinje et al., Nucl. Acids. Res. 14:4683-4690 (1986)). Moreover, it is also recognized that, in some cases, cleavage of a signal sequence from a secreted polypeptide is not entirely uniform, resulting in more than one secreted species. These mature polypeptides, where the signal peptide is cleaved within no more than about 5 amino acids on either side of the C-terminal boundary of the signal peptide as identified herein, and the polynucleotides encoding them, are contemplated by the present invention.

"PRO polypeptide variant" means an active PRO polypeptide as defined above or below having at least about 80% amino acid sequence identity with a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Such PRO polypeptide variants include, for instance, PRO polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of the full-length native amino acid sequence. Ordinarily, a PRO polypeptide variant will have at least about 80% amino acid sequence identity, preferably at least about 81% amino acid sequence identity, more preferably at least about 82% amino acid sequence identity, more preferably at least about 83% amino acid sequence identity, more preferably at least about 84% amino acid sequence identity, more preferably at least about 85% amino acid sequence identity, more preferably at least about 86% amino acid sequence identity, more preferably at least about 87% amino acid sequence identity, more preferably at least about 88% amino acid sequence identity, more preferably at least about 89% amino acid sequence identity, more preferably at least about 90% amino acid sequence identity, more preferably at least about 91% amino acid sequence identity, more preferably at least about 92% amino acid sequence identity, more preferably at least about 93% amino acid sequence identity, more preferably at least about 94% amino acid sequence identity, more preferably at least about 95% amino acid sequence identity, more preferably at least about 96% amino acid sequence identity, more preferably at least about 97% amino acid sequence identity, more preferably at least about 98% amino acid sequence identity and most preferably at least about 99% amino acid sequence identity with a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, PRO variant polypeptides are at least about 10 amino acids in length, often at least about 20 amino acids in length, more often at least about 30 amino acids in length, more often at least about 40 amino acids in length, more often at least about 50 amino acids in length, more often at least about 60 amino acids in length, more often at least about 70 amino acids in length, more often at least about 80 amino acids in length, more often at least about 90 amino acids in length, more often at least about 100 amino acids in length, more often at least about 150 amino acids in length, more often at least about 200 amino acids in length, more often at least about 300 amino acids in length, or more.

"Percent (%) amino acid sequence identity" with respect to the PRO polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific PRO polypeptide sequence, after aligning the sequences and introducing gaps,

D. Preparation of PRO

The description below relates primarily to production of PRO by culturing cells transformed or transfected with a vector containing PRO nucleic acid. It is, of course, contemplated that alternative methods, which are well known in the art, may be employed to prepare PRO. For instance, the PRO sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., Solid-Phase Peptide Synthesis, W.H. Freeman Co., San Francisco, CA (1969); Merrifield, J. Am. Chem. Soc., 85:2149-2154 (1963)]. *In vitro* protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, CA) using manufacturer's instructions. Various portions of the PRO may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length PRO.

1. Isolation of DNA Encoding PRO

DNA encoding PRO may be obtained from a cDNA library prepared from tissue believed to possess the PRO mRNA and to express it at a detectable level. Accordingly, human PRO DNA can be conveniently obtained from a cDNA library prepared from human tissue, such as described in the Examples. The PRO-encoding gene may also be obtained from a genomic library or by known synthetic procedures (e.g., automated nucleic acid synthesis).

Libraries can be screened with probes (such as antibodies to the PRO or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding PRO is to use PCR methodology [Sambrook et al., supra; Dieffenbach et al., PCR Primer: A Laboratory Manual (Cold Spring Harbor Laboratory Press, 1995)].

The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like ³²P-labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., supra.

Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined using methods known in the art and as described herein.

Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook et al., supra, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

2. Selection and Transformation of Host Cells

Host cells are transfected or transformed with expression or cloning vectors described herein for PRO production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in Mammalian Cell Biotechnology: a Practical Approach, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., supra.

Methods of eukaryotic cell transfection and prokaryotic cell transformation are known to the ordinarily skilled artisan, for example, CaCl_2 , CaPO_4 , liposome-mediated and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., supra, or electroporation is generally used for prokaryotes. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., Gene, 23:315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transfections have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., J. Bact., 130:946 (1977) and Hsiao et al., Proc. Natl. Acad. Sci. (USA), 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown et al., Methods in Enzymology, 185:527-537 (1990) and Mansour et al., Nature, 336:348-352 (1988).

Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published 12 April 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype *tonA*; *E. coli* W3110 strain 9E4, which has the complete genotype *tonA ptr3*; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT kan^r*; *E. coli* W3110 strain 37D6, which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7*

ilvG kan^r; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant *degP* deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Patent No. 4,946,783 issued 7 August 1990. Alternatively, *in vitro* methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for PRO-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, Nature, 290: 140 [1981]; EP 139,383 published 2 May 1985); *Kluyveromyces* hosts (U.S. Patent No. 4,943,529; Fleer et al., Bio/Technology, 9:968-975 (1991)) such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt et al., J. Bacteriol., 737 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilae* (ATCC 36,906; Van den Berg et al., Bio/Technology, 8:135 (1990)), *K. thermotolerans*, and *K. marxianus*; *Yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna et al., J. Basic Microbiol., 28:265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case et al., Proc. Natl. Acad. Sci. USA, 76:5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 October 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance et al., Biochem. Biophys. Res. Commun., 112:284-289 [1983]; Tilburn et al., Gene, 26:205-221 [1983]; Yelton et al., Proc. Natl. Acad. Sci. USA, 81: 1470-1474 [1984]) and *A. niger* (Kelly and Hynes, EMBO J., 4:475-479 [1985]). Methylophilic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, The Biochemistry of Methylophilic Yeasts, 269 (1982).

Suitable host cells for the expression of glycosylated PRO are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful mammalian host cell lines include Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen. Virol., 36:59 (1977)); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod., 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

3. Selection and Use of a Replicable Vector

The nucleic acid (e.g., cDNA or genomic DNA) encoding PRO may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally

include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

5 The PRO may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the PRO-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g.,
10 the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral
15 secretory leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for
20 cloning vectors in mammalian cells.

Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

25 An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the PRO-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., Proc. Natl. Acad. Sci. USA, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb et al., Nature,
30 282:39 (1979); Kingsman et al., Gene, 7:141 (1979); Tschemper et al., Gene, 10:157 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, Genetics, 85:12 (1977)].

Expression and cloning vectors usually contain a promoter operably linked to the PRO-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well
35 known. Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems [Chang et al., Nature, 275:615 (1978); Goeddel et al., Nature, 281:544 (1979)], alkaline phosphatase, a tryptophan (*trp*) promoter system [Goeddel, Nucleic Acids Res., 8:4057 (1980); EP 36,776], and hybrid

promoters such as the tac promoter [deBoer et al., Proc. Natl. Acad. Sci. USA, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding PRO.

5 Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., J. Biol. Chem., 255:2073 (1980)] or other glycolytic enzymes [Hess et al., J. Adv. Enzyme Reg., 7:149 (1968); Holland, Biochemistry, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

10 Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

15 PRO transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

20 Transcription of a DNA encoding the PRO by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the PRO coding sequence, but is preferably located at a site 5' from the promoter.

25 Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding PRO.

30 Still other methods, vectors, and host cells suitable for adaptation to the synthesis of PRO in recombinant vertebrate cell culture are described in Gething et al., Nature, 293:620-625 (1981); Mantei et al., Nature, 281:40-46 (1979); EP 117,060; and EP 117,058.

4. Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)], dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence PRO polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to PRO DNA and encoding a specific antibody epitope.

5. Purification of Polypeptide

Forms of PRO may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of PRO can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

It may be desired to purify PRO from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the PRO. Various methods of protein purification may be employed and such methods are known in the art and described for example in Deutscher, Methods in Enzymology, 182 (1990); Scopes, Protein Purification: Principles and Practice, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular PRO produced.

E. Uses for PRO

Nucleotide sequences (or their complement) encoding PRO have various applications in the art of molecular biology, including uses as hybridization probes, in chromosome and gene mapping and in the generation of anti-sense RNA and DNA. PRO nucleic acid will also be useful for the preparation of PRO polypeptides by the recombinant techniques described herein.

PRO330 [herein designated as DNA40603-1232] (SEQ ID NO:331) and the derived protein sequence for PRO330.

The entire nucleotide sequence of DNA40603-1232 is shown in Figure 115 (SEQ ID NO:331). Clone DNA40603-1232 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 167-169 and ending at the stop codon at nucleotide positions 1766-1768 (Figure 115). The predicted polypeptide precursor is 533 amino acids long (Figure 116). Clone DNA40603-1232 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209486 on November 21, 1997.

Analysis of the amino acid sequence of the full-length PRO330 polypeptide suggests that portions of it possess significant homology to the mouse prolyl 4-hydroxylase alpha subunit protein, thereby indicating that PRO330 may be a novel prolyl 4-hydroxylase alpha subunit polypeptide.

EXAMPLE 49: Isolation of cDNA Clones Encoding Human PRO310

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA40553. Based on the DNA40553 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO310.

Forward and reverse PCR primers were synthesized:

forward PCR primer 1 5'-TCCCCAAGCCGTTCTAGACGCGG-3' (SEQ ID NO:342)

forward PCR primer 2 5'-CTGGTTCTTCCTTGCACG-3' (SEQ ID NO:343)

reverse PCR primer 5'-GCCCAAATGCCCTAAGGCGGTATACCCC-3' (SEQ ID NO:344)

Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus sequence which had the following nucleotide sequence

hybridization probe

5'-GGGTGTGATGCTTGAAGCATTTTCTGTGCTTTGATCACTATGCTAGGAC-3' (SEQ ID NO:345)

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO310 gene using the probe oligonucleotide and one of the PCR primers.

RNA for construction of the cDNA libraries was isolated from human fetal liver tissue.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO310 [herein designated as DNA43046-1225 (SEQ ID NO:340) and the derived protein sequence for PRO310 (SEQ ID NO:341).

The entire nucleotide sequence of DNA43046-1225 is shown in Figure 119 (SEQ ID NO:340). Clone DNA43046-1225 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 81-83 and ending at the stop codon at nucleotide positions 1035-1037 (Figure 119). The predicted polypeptide precursor is 318 amino acids long (Figure 120) and has a calculated molecular weight of approximately 36,382 daltons. Clone DNA43046-1225 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209484.

Analysis of the amino acid sequence of the full-length PRO310 polypeptide suggests that portions of it possess homology to *C. elegans* proteins and to fringe, thereby indicating that PRO310 may be involved in development.

EXAMPLE 50: Isolation of cDNA clones Encoding Human PRO339

5 An expressed sequence tag (EST) DNA database (LIFESEQ™, Incyte Pharmaceuticals, Palo Alto, CA) was searched and ESTs were identified. An assembly of Incyte clones and a consensus sequence was formed using phrap as described in Example 1 above.

Forward and reverse PCR primers were synthesized based upon the assembly-created consensus sequence:

- 10 forward PCR primer 1 5'-GGGATGCAGGTGGTGTCTCATGGGG-3' (SEQ ID NO:346)
forward PCR primer 2 5'-CCCTCATGTACCGGCTCC-3' (SEQ ID NO:347)
forward PCR primer 3 5'-GTGTGACACAGCGTGGGC-3' (SEQ ID NO:43)
forward PCR primer 4 5'-GACCGGCAGGCTTCTGCG-3' (SEQ ID NO:44)
reverse PCR primer 1 5'-CAGCAGCTTCAGCCACCAGGAGTGG-3' (SEQ ID NO:45)
15 reverse PCR primer 2 5'-CTGAGCCGTGGGCTGCAGTCTCGC-3' (SEQ ID NO:46)

Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus sequence which had the following nucleotide sequence

hybridization probe

5'-CCGACTACGACTGGTTCTTCATCATGCAGGATGACACATATGTGC-3' (SEQ ID NO:47)

20 In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified above. A positive library was then used to isolate clones encoding the PRO339 gene using the probe oligonucleotide and one of the PCR primers.

RNA for construction of the cDNA libraries was isolated from human fetal liver tissue.

25 A cDNA clone was sequenced in entirety. The entire nucleotide sequence of DNA43466-1225 is shown in Figure 117 (SEQ ID NO:338). Clone DNA43466-1225 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 333-335 and ending at the stop codon found at nucleotide positions 2649-2651 (Figure 117; SEQ ID NO:338). The predicted polypeptide precursor is 772 amino acids long and has a calculated molecular weight of approximately 86,226 daltons. Clone DNA43466-1225 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209490.

30 Based on a BLAST and FastA sequence alignment analysis (using the ALIGN computer program) of the full-length sequence, PRO339 has homology to *C. elegans* proteins and collagen-like polymer sequences as well as to fringe, thereby indicating that PRO339 may be involved in development or tissue growth.

EXAMPLE 51: Isolation of cDNA Clones Encoding Human PRO244

35 A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. Based on this consensus sequence, oligonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length

FIGURE 120

MLSESSSFLKGVMLGSIFCALITMLGHIRIGHGNRMHHHEHHHLQAPNKEDILKISEDERME
LSKSFRVYCIILVKPKDVSLWAAVKETWTKHCDKAEFFSSENVKVFESINMDTNDMWLMMRK
AYKYAFDKYRDQYNWFFLARPTTFAI IENLKYFLLKKDPSQPFYLGHTIKSGDLEYVGMEGG
IVLSVESMKRLNSLLNIPEKCPEQGGMWKISEDKQLAVCLKYAGVFAENAEDADGKDVFNT
KSVGLSIKEAMTYHPNQVVEGCCSDMAVTFNGLTPNQMHVMMYGVYRLRAFGHIFNDALVFL
PPNGSDND

Signal sequence:

amino acids 1-33

N-glycosylation site.

amino acids 121-125, 342-346

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 319-323, 464-468

Casein kinase II phosphorylation site.

amino acids 64-132, 150-154, 322-326, 331-335, 368-372, 385-389,
399-403, 409-413, 473-477, 729-733, 748-752

Tyrosine kinase phosphorylation site.

amino acids 736-743

N-myristoylation site.

amino acids 19-25, 23-29, 136-142, 397-403, 441-447, 544-550,
558-564, 651-657, 657-663, 672-672

Prokaryotic membrane lipoprotein lipid attachment site.

amino acids 14-25

Cell attachment sequence.

amino acids 247-250